



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

If R , r , r_1 , r_2 , and r_3 be, respectively, the radii of the circumscribed, inscribed, and escribed circles of a \triangle , prove $r_1 + r_2 + r_3 - r = 4R$.

41. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Find the length (x) of a rectangular parallelepiped $b=5$ ft. and $h=3$ ft., which can be *diagonally inscribed* in a similar parallelepiped $L=83$ ft., $B=64$ ft., and $H=50$ ft.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

26. Proposed by Professor J. F. W. SCHEFFER, M. A., Hagerstown, Maryland.

According to Bessel, the ratio of the squares of the polar diameter of the earth to that of the equatorial diameter, is .9933254. Find what *latitude* the angle made by a body falling to the earth, with a perpendicular to the surface, is greatest. Find, also, this maximum angle.

Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Let ϕ = the required geographical latitude, and ϕ' = the geocentric latitude of the same place; then *Chauvenet's Spherical and Practical Astronomy*, Vol. I., p. 98, we deduce $\phi' = \tan^{-1}[(b^2/a^2)\tan\phi] = \tan^{-1}[(1-e^2)\tan\phi]$.

$$\therefore (\phi - \phi') = \phi - \tan^{-1}[(1-e^2)\tan\phi], = \text{a Maximum.}$$

$$\therefore \frac{d(\phi - \phi')}{d\phi} = 1 - \frac{(1-e^2)(1+\tan^2\phi)}{1+(1+e^2)^2\tan^2\phi} = 0.$$

$$\therefore \phi = \tan^{-1} \left[\sqrt{\left(\frac{1}{1-e^2} \right)} \right] = \tan^{-1}(1.0033541) = 45^\circ 5' 45''.32,$$

$$\text{and } \phi' = \tan^{-1}(.9966571) = 44^\circ 54' 14''.67.$$

Hence $(\phi - \phi') = 11' 30''.65$; and this result is found in the already-named *Manual of Astronomy*, Vol. II.; third Table, p. 577.

II. Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.


Not taking into account the eastward deviation due to the rotation of the earth we can proceed as follows:

Let HEC be the direction the body falls, FEG the perpendicular to the earth's surface at E , DEK the tangent to the meridian at E , $CA=a$,

$CB=b$, $\angle ECA=\theta$, $\angle DCE=\angle CEG=\phi$, $\angle EKX=\beta$, co-ordinates of $E=(x,y)$.

Then $\tan \beta = -\frac{b^2 x}{a^2 y}$, $\tan \theta = \frac{y}{x}$, also
 $\beta = 90^\circ + \theta + \phi$, $\tan \beta = \tan (90^\circ + \theta + \phi)$
 $= -\cot (\theta + \phi)$.

$$\begin{aligned}\therefore \frac{b^2 x}{a^2 y} &= \cot (\theta + \phi) = \frac{\cot \theta \cot \phi - 1}{\cot \theta + \cot \phi} \\ &= \frac{\frac{x}{y} \cot \phi - 1}{\frac{x}{y} + \cot \phi}.\end{aligned}$$



$$\therefore \tan \phi = \frac{a^2 - b^2}{a^2 b^2} xy = \text{maximum} \dots (1). \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \dots (2).$$

The first differentials of (1) and (2) give $b^2 x^2 = a^2 y^2$.

$$\therefore x = \frac{a}{\sqrt{2}}, y = \frac{b}{\sqrt{2}}, \therefore \tan \theta = \frac{b}{a} = (.9933254)^2 = .996659, \therefore \theta = 44^\circ 54'$$

$$14'' .9 = \text{the latitude} \quad \tan \phi = \frac{a^2 - b^2}{2ab} = \frac{.0066746}{1.993318} = .003348, \therefore \phi = 1' 30'' .5$$

= maximum angle made with the perpendicular.

Also solved by Professor C. W. M. Black, and the Proposer.

PROBLEMS.

36. Proposed by H. C. WHITAKER, B. Sc., M. E., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

A cube is revolved on its diagonal as an axis. Define the figure described and calculate its volume.

37. Proposed by J. A. CALDERHEAD, Superintendent of Schools, Lima, Ohio.

A man ties two mules—one to the outside of a circular wall, the other to the inside. If the lengths of the ropes of each is one-fourth the circumference of the wall, and both together can graze over one acre of ground: find the circumference of the wall.